

⑫

EUROPEAN PATENT APPLICATION

⑲ Application number: 86117748.3

⑳ Date of filing: 19.12.86

⑤① Int. Cl.⁴: **C 07 C 103/46**
C 07 C 103/84, C 07 C 103/7-
37
C 07 C 147/00, C 07 C 157/0-
7
C 07 D 257/04, A 61 K 31/16-
5
A 61 K 31/41

③① Priority: 20.12.85 US 811567

④③ Date of publication of application:
24.06.87 Bulletin 87/26

④④ Designated Contracting States:
AT BE CH DE ES FR GB GR IT LI LU NL SE

⑦① Applicant: **WARNER-LAMBERT COMPANY**
201 Tabor Road
Morris Plains New Jersey 07950(US)

⑦② Inventor: **Rafferty, Michael F.**
59 Clear Lake Manor Drive East
North Branford Connecticut 06471(US)

⑦③ Inventor: **Johnson, Graham**
1130 Bandera Drive
Ann Arbor Michigan 48103(US)

⑦④ Representative: **Mansmann, Ivo**
c/o Gödecke AG - Patentabteilung Postfach 569 -
Mooswaldallee 1-9
D-7800 Freiburg(DE)

⑤④ **Benzolc acid and benzoic acid ester derivatives having anti-inflammatory and analgesic activity.**

⑤⑦ Novel benzoic acid or benzoic acid ester derivatives, pharmaceutical compositions and methods of use thereof are the present invention. Utility is for the treatment of arthritis, asthma, Raynaud's disease, inflammatory bowel disorders, trigeminal or herpetic neuralgia, inflammatory eye disorders, psoriasis, dental pain, and headaches, particularly vascular headache, such as migraine, cluster, mixed vascular syndromes, as well as nonvascular, tension headaches.

EP 0 226 223 A2

BEST AVAILABLE COPY

The present invention is novel compounds, which are derivatives of benzoic acid and benzoic acid esters--having antiinflammatory activity for the treatment of arthritis, asthma, Raynaud's disease, inflammatory bowel disorders, trigeminal or herpetic neuralgia, inflammatory eye disorders, psoriasis, and/or having analgesic activity for the treatment of dental pain and headache, particularly vascular headache, such as migraine, cluster, and mixed vascular syndromes, as well as nonvascular, tension headache. Thus, the present invention is also a pharmaceutical composition comprising the novel compounds together with a pharmaceutically acceptable carrier or methods of use of such compounds comprising the preparation of pharmaceuticals for treatment of the above noted conditions.


Among known compounds are benzoic acid derivations in which the derivative is limited to a substituent having a (naphthoxy)isobutyramido containing group and for which compounds an antiphlogistic activity is disclosed. See U.S. Patent No. 4,183,954. Additionally O. Exner, et al. discloses N-(4-carboxybenzyl)acetamide in "Quantitative Evaluation of the Inductive Effect," Coll. Czech. Chem. Commun. 27, 2299 (1962). But no teaching to activity or utility for the compound is indicated by Exner, et al.

Compounds related to capsaicin are disclosed in a series of patents. The compounds are thus not benzoic acid derivatives but have various amido, sulfonylamido or amidosulfonyl and thioamido linkages in combination with a benzyl or a benzyl analog moiety. Such compounds are found in U.S. Patent No. 4,313,958, that

claims the use of capsaicin; U.S. Patent No. 4,460,602;
 U.S. Patent No. 4,401,663 and U.S. Patent No. 4,443,477;
 European Patent Application 0,132,113; U.S. Patent No.
 4,424,203; European Patent Application 0,132,114;
 5 European Patent Application 0,132,346 and European Patent
 Application 0,132,115 as well as European Patent Application
 0,149,544 and 0,149,545. Of these European Patent
 Applications 0,132,115; 0,132,346; 0,132,114; 0,132,115
 0,149,544 and 0,149,545 include a short chain acyl group
 10 on the benzyl moiety. U.S. Patent No. 3,992,540 discloses
 3-guino-line-carboxamides. Analgesia is disclosed as an
 activity for the compounds of the references. However,
 none of the references teach the compounds having the
 moieties such as benzoic acid moieties and their substituents,
 15 or particularly the combination of moieties, of the present
 invention.

The novel compounds of the present invention have
 the following structural formula I:


20




wherein:

(a) R_1 is tetrazolyl or COOR' wherein R' is H or lower
 alkyl of 1 to 4 carbons, inclusive;


(b) B is




0226223



- (c) X and Y are independently H or lower alkyl of 1 to 4 carbons, inclusive;
- 5 (d) R_2 is alkylene, alkenylene, alkynylene branched or linear chains of 1 to 11 carbons, inclusive;
- (e) Q is CH_3 , COOH , Br, NH_2 , H, imidazolyl, cyclohexyl,



10 or




- wherein R_3 and R_4 are hydrogen or a C_1 to C_4 alkyl- or C_1 to C_4 alkoxy group, and nontoxic, pharmaceutically acceptable base or acid addition salts thereof, with the proviso that when B is (B_1) and Q is H, then R_2 is not methylene and with the
- 15 further proviso that when B is (B_2) then Q cannot be CH_3 , H or phenyl.

The term "lower alkyl of 1 to 4 carbons" means a straight or branched hydrocarbon chain up to 4 carbon atoms such as methyl, ethyl, propyl, isopropyl, butyl, isobutyl, secondary butyl or tertiary butyl.

The terms alkylene, alkenylene and alkynylene are divalent hydrocarbon straight or branched chains containing one or more single, double or triple carbon to carbon bonds, respectively.

- 5 In the preferred embodiments of the present invention R_1 is COOH as shown in the following formula (II):



II

- wherein X, Y, B, R_2 and Q are all as defined above. More preferred embodiments of the present invention are
 10 compounds of formula II wherein B is B_1 and X, Y, R_2 and Q are as defined above. The most preferred embodiment of the present invention is the compound N-(4-carboxybenzyl)nonanamide. The preferred method of use is for treating headaches, particularly migraines.


- 15 Examples of suitable acids for the preparation of the acid addition salts are inorganic acids such as hydrochloric acid, hydrobromic acid, sulfuric acid, phosphoric acid, and the like, and organic acids such as acetic acid, benzoic acid, tartaric acid, fumaric
 20 acid, succinic acid, maleic acid, arginine acid, lactic acid, tartaric acid, and sulfonic acids such as methansulfonic acid, ethansulfonic acid, benzenesulfonic acid or p-toluenesulfonic acid.

- The base salts of the present inventions include
 25 those safe for topical or systemic administration, such

0226223

as sodium, potassium, calcium, magnesium, and ammonium salts or the like.


Generally, the preparation of the compounds of the present invention is represented by the following
5 scheme:



wherein R₁, X, Y, B, R₂, Q are as defined above and Hal is chloro, bromo, or iodo, but preferably chloro.


The preparation uses standard synthetic
10 techniques used in the examples or analogous to those used in the examples hereinafter. The starting materials for the preparation are readily available, known or can be prepared by known methods.

The compositions containing the compounds of the
15 formula (I')




wherein:


- (a) R₁ is tetrazolyl or COOR' wherein R' is H or lower alkyl of 1 to 4 carbons, inclusive;
20 (b) B is



0226223



- (c) X and Y are independently H or lower alkyl of 1 to 4 carbons, inclusive;
- 5 (d) R_2 is alkylene, alkenylene, alkynylene branched or linear chains of 1 to 11 carbons, inclusive;
- (e) Q is CH_3 , COOH , Br, NH_2 , H, imidazolyl, cyclohexyl,



10 or



wherein R_3 and R_4 are hydrogen or a C_1 to C_4 alkyl or a C_1 to C_4 alkoxy group.

and nontoxic, pharmaceutically acceptable base or acid addition salts thereof, are comprised of an analgesic or antiinflammatory effective amount of a compound of

15 formula I' as defined above or their pharmaceutically acceptable base or acid addition salts and a pharmaceutically acceptable carrier. Such

0226223

compositions may be one of a broad range of known forms for topical or systemic administration.

The compounds I' may be used for the treatment in mammals, particularly in humans, of various conditions
5 such as enumerated above either for diseases known as inflammatory or for pain. An ordinarily skilled physician would recognize such conditions. The compounds of formula I are active in animal tests which are generally recognized as predictive for
10 antiinflammatory or analgesic activity. Regardless of the route of administration selected, the compounds of the present invention are formulated into pharmaceutically acceptable dosage forms by conventional methods known to the pharmaceutical art.
15 In general a preferred method of administration is, however, by oral dosage forms.

The compounds can be administered in such unit oral dosage forms as tablets, capsules, pills, powders, or granules. They may also be administered rectally or
20 vaginally in such forms as suppositories or bougies. They may also be introduced parenterally, (e.g., subcutaneously, intravenously, or intramuscularly), using forms known to the pharmaceutical art.

An effective but nontoxic amount of the compound
25 of formula I or the salts thereof is employed in treatment. The dosage regimen for treating inflammation or pain by the compounds of formula I and their salts as described above is selected in accordance with a variety of factors including the
30 type, age, weight, sex, and medical condition of the subject, the severity of the inflammation or pain, the route of administration and the particular compound employed. Determination of the proper dosage for a particular situation is within the skill of the art.

0226223

Generally, treatment is initiated with smaller dosages which are less than the optimum dose of the compound. Thereafter the dosage is increased by small increments until the optimum effect under the circumstances is reached. For convenience, the total daily dosage may be divided and administered in portions during the day if desired.

Initial dosages of the compounds of the invention are ordinarily in the area of 1 mg/kg up to at least 100 mg/kg per dose orally, preferably 30 to 100 mg/kg orally are given. Each dose is given from one to four times daily or as needed. When other forms of administration are employed equivalent doses are administered.

An illustrative example of the activity for use as described above for the novel compounds of the present invention is an ED_{50} of 33.03 mg/kg for the compound of Example 1 described in the following material when administered in a test based on that of Koster et al. [Fed. Proc., Vol. 18 (1959), p. 412] in which the peritoneal injection of acetic acid to mice provokes repeated stretching and twisting movements which persist for more than 6 hrs. Analgesics prevent or suppress these syndromes which are considered to be an exteriorization of a diffuse abdominal pain. A 1% solution of acetic acid in water is used at a dose of 0.01 ml/g or 100 mg/kg of acetic acid to release the syndrome.

The Example 1 compound is subcutaneously administered 30 minutes before the acetic acid injection and the mice are fasted 24 hrs before the start of the test. The stretching for the mice is observed and totaled for each mouse in a 15 minute observation period starting just after the acetic acid

0226223

injection. The results are expressed as mg/kg which amount produces the desired inhibition of stretching or "writhing" in 50 percent of a population.

5 Additionally, the same Example 1 compound was effective at a dose of 100 mg/kg administered i.p. in reducing the inflammatory response to an injection of carrageenan into the rat foot pad. This is a commonly employed standard assay for the identification of antiinflammatory activity, based on the method
10 described by Winter et al. (Proc. Soc. Exptl. Biol. N.Y. vol 111 (1962), p. 544).

The following Examples will further illustrate the invention, without limiting it thereto.

EXAMPLE 1

N-(4-Carboxybenzyl)nonanamide

Methyl 4-(aminomethyl)benzoate hydrochloride (6.0 g, 0.03 moles) is treated with 50 mL 1 N NaOH and the mixture extracted with ether (3 x 50 mL). The combined ether extracts are dried with anhydrous potassium carbonate and evaporated to leave the amine base as a white solid. This residue is dissolved in 100 mL methylene chloride, to which is added 2.85 g pyridine. Nonanoyl chloride (6.36 g, 0.036 moles) in 10 mL methylene chloride is then added dropwise to the mixture with stirring over a 5 min period. The thick pasty mass which formed after a few minutes is stirred at room temperature for 45 min, at which time 30 mL saturated sodium bicarbonate is carefully added. The mixture is stirred vigorously for 15 min, after which the layers are separated and the organic layer extracted with 2 N HCl (30 mL), and dried over anhydrous sodium sulfate. Evaporation of the solvent leaves a waxy residue which is crystallized from isopropyl ether as colorless plates, m.p. 94.5-95.5°C.

The crystalline product (2.0 g) is dissolved in tetrahydrofuran (30 mL) to which is added 1 N NaOH (10 mL). The heterogeneous mixture is stirred overnight at room temperature. The resulting clear solution is made acidic by addition of 2 N HCl, and the mixture partitioned between chloroform (250 mL) and water (200 mL). The chloroform layer is dried over sodium sulfate and evaporated to leave a waxy residue, which is crystallized as colorless needles from methanol/water. A yield of 1.36 g, of the desired product N-(4-carboxybenzyl)nonanamide is obtained. M.p. 178-179.5°C.

In a procedure analogous to that described in Example 1 above but using the appropriate acid chloride the following compounds are prepared.

0226223

EXAMPLE 2

N-(4-Carboxybenzyl)decanamide sodium salt, m.p. 250°C.

EXAMPLE 3

N-(4-Carboxybenzyl)heptanamide, m.p. 179-180°C.

5 EXAMPLE 4

N-(4-Carboxybenzyl)octanamide, m.p. 180°C.

EXAMPLE 5

N-(4-Carboxybenzyl)phenylacetamide, m.p. 220-221°C.

EXAMPLE 6

10 N-(4-Carboxybenzyl)-4-hydroxy-3-methoxycinnamamide,
m.p. 233-234°C.

EXAMPLE 7

N-(4-Carboxybenzyl)-4-phenylbutyramide

15 4-(Aminomethyl)benzoic acid (3.6 g) is suspended
in methylene chloride (100 mL), to which 15 mL
triethylamine is added. Chlorotrimethylsilane (10 mL)
is then added and the mixture allowed to stir at room
temperature for 1 hr. The mixture is then cooled in an
ice bath and 4-phenylbutyryl chloride (5.3 g) in
20 methylene chloride (10 mL) is added dropwise and the
resulting mixture stirred for 30 min at 0°C, followed
by an additional 3 hrs at ambient temperature. The
mixture is treated with 75 mL 1 N HCl, after which the
organic layer is separated and extracted with 1 N HCl.

0226223

The precipitate which had formed is recovered by filtration and recrystallized two times from methanol/1 N HCl to give pure N-(4-carboxybenzyl)-4-phenylbutyramide. M.p. 178-179°C.

- 5 A procedure analogous to that described in Example 7 using the appropriate starting material produces the following compound:

EXAMPLE 8

N-(4-Carboxybenzyl)undecanamide, m.p. 179-180°C.

10 EXAMPLE 9

N-(4-Carboxybenzyl)-(2-naphthoxy)acetamide

- (2-Naphthoxy)acetic acid (8.5 g) is suspended in 160 mL methylene chloride and treated with 1,1'-carbonyldiimidazole (6.8 g) which is added in small portions. After stirring for 3 hrs at room temperature under a nitrogen atmosphere, the mixture is added dropwise to a previously prepared solution of alpha-amino-p-toluic acid (1.4 g), chlorotrimethylsilane (10 mL), and triethylamine (11 mL) in 250 mL methylene chloride at 0°C. The final mixture is stirred at room temperature under a nitrogen atmosphere overnight. The mixture is combined with 200 mL 1 N HCl and shaken, after which the resultant precipitate is collected by suction filtration. The precipitate is recrystallized from methanol/2N HCl, m.p. 188-189°C as N-(4-carboxybenzyl)-(2-naphthoxy)acetamide.
- 15
20
25

EXAMPLE 10

N-(4-Carboxybenzyl)cinnamamide

- 1,1'-Carbonyldiimidazole (3.71 g) is added to an ice-cold stirred solution of cinnamic acid (3.13 g) in 30 mL tetrahydrofuran. After stirring for an additional 1 hr, methyl 4-(aminomethyl)benzoate
- 30

0226223

hydrochloride (4.7 g) and triethylamine (3.23 mL) are added and the final mixture stirred while immersed in an ice water bath for an additional 30 min, followed by stirring overnight at room temperature. After removal
5 of the solvent by rotary evaporation, the residue is taken up in chloroform (250 mL) and extracted with water (200 mL), 1N HCl (3 x 50 mL), water (100 mL), saturated sodium bicarbonate (100 mL), brine (100 mL) and the final chloroform layer dried over anhydrous
10 magnesium sulfate. After evaporation of the solvent in vacuo, the crude product (2.74 g) is suspended in 100 mL tetrahydrofuran and 20 mL 2 N NaOH and the mixture stirred at room temperature overnight. The mixture is then made acidic by addition of excess 1 N HCl, and the
15 precipitate recovered by filtration. After washing the filter cake with water and pressing to remove as much water as possible, the white solid is crystallized from methanol/2N HCl as N-(4-carboxybenzyl)cinnamamide, m.p. 238-239°C.

20 A procedure analogous to that described in Example 10 using an appropriate starting material produces the following compound:

EXAMPLE 11

N-(4-Carboxybenzyl)cyclohexylacetamide, m.p. 223-
25 224°C.

EXAMPLE 12

N-(4-Carboxybenzyl)butyramide

4-(Aminomethyl)benzoic acid (3.0 g) is suspended in pyridine (15 mL) and the mixture cooled in an ice
30 water bath. Butyric anhydride (9.2 mL) is added dropwise to the stirred suspension over a 20 min period. The ice bath is removed and the mixture stirred at room temperature overnight. The mixture is poured into 150 mL ice water and made acidic (pH 1.5)

0226223

by addition of concentrated HCl. The precipitate is recovered by suction filtration and recrystallized from ethyl acetate to produce N-(4-carboxybenzyl)butyramide, m.p. 186.5-187.5°C.

5 EXAMPLE 13

N-(4-Carboxybenzyl)hexanamide

Hexanoic acid (3.06 g) in 20 mL acetonitrile is treated with N-methylmorpholine (2.9 mL) and the mixture cooled to -20°C with stirring. Ethyl
10 chloroformate (2.8 mL) is then added dropwise, keeping the temperature below or at -20°C. After stirring for an additional 40 min at that temperature, the solution is transferred to a cooled (-15°C) solution of alpha-amino-p-toluic acid (2.0 g), triethylamine (15 mL), a
15 chlorotrimethylsilane (5.0 mL) in methylene chloride (60 mL; prepared as described in Example 7). After the addition is complete, the mixture is stirred at 5°C for 4 hrs, followed by overnight stirring at room temperature. After removal of the solvents by
20 evaporation, the residue is redissolved in methylene chloride (100 mL) and extracted with 1N HCl (2x50 mL) and brine (2x50 mL). The organic layer is dried over magnesium sulfate and evaporated, leaving N-(4-carboxybenzyl)hexanamide as an off-white solid. The N-
25 (4-carboxybenzyl)hexanamide product is crystallized from methanol/2N HCl, m.p. 178-179°C.

EXAMPLE 14

N-1-(1-(4-Carboxyphenyl)ethyl)nonanamide

Step 1 4-(1-aminoethyl)benzoic acid

30 4-Acetylbenzoic acid (4.1 g) is dissolved in 50 mL ammonia-saturated methanol. Raney nickel catalyst (1.5 g; activity grade III) is then added and the mixture reduced under hydrogen atmosphere (4750 psi) at 80°C for 17 hrs. After removal of the catalyst by

0226223

suction filtration, the filtrate is evaporated and the residue dissolved in H_2O . The solution is passed through a 2.5x15 cm column of Dowex 50X8-400 resin (H^+ form) and eluted with 1N NH_4OH . Evaporation of the eluate leaves a residue (2.9 g) which is recrystallized from H_2O /acetone and characterized as 4-(1-aminoethyl)benzoic acid, m.p. $> 300^\circ C$.

Step 2 N-1-(1-(4-Carboxyphenyl)ethyl)nonanamide

The 4-(1-aminoethyl)benzoic acid as prepared above in Step 1 (1.5 g) is suspended in 30 mL methylene chloride containing 2.13 g pyridine and cooled to $0^\circ C$. Nonanoyl chloride (1.7 g) is dissolved in 5 mL methylene chloride and added dropwise with stirring to the cooled solution. After allowing the mixture to warm to room temperature, the mixture is allowed to stir an additional 2 hrs. Treatment with 1 N HCl (40 mL) produces a solid residue at the interface of the two liquid phases, which is recovered by filtration and recrystallized from methanol/water as N-1-(1-(4-carboxyphenyl)ethyl)nonanamide, m.p. $178-180^\circ C$.

EXAMPLE 15

N-(4-carboxybenzyl)-N-methylnonanamide

Step 1 4-(methylaminomethyl)benzoic acid hydrochloride

4-Carboxybenzaldehyde (10 g) is dissolved in 50 mL aqueous methylamine (30%). Raney nickel (5 g) is added and the mixture treated with hydrogen at 1500 psi and $100^\circ C$ for 17 hrs. Removal of the catalyst by suction filtration and evaporation of the filtrate left a solid residue, which is redissolved in 2 N HCl (50 mL). The solution is extracted with ethyl acetate (50 mL) and chloroform (50 mL) and the resultant aqueous layer evaporated to dryness. The residue is vacuum dried at $60^\circ C$ for 4 hrs and recrystallized from methanol/ethyl acetate to yield 7.65 g 4-(methylaminomethyl) benzoic acid hydrochloride, m.p. $255-261^\circ C$.

5 Step 2 N-(4-Carboxybenzyl-N-methylnonanamide)

The 4-(methyaminomethyl)benzoic acid hydrochloride prepared in Step 1 above (2.0 g) is suspended in 5 mL pyridine and cooled in ice water. Nonanoyl chloride (1.8 g) is added dropwise with
10 stirring, and the final solution allowed to stir at room temperature for 18 hrs. The clear solution is treated with 2 N HCl (15 mL), and partitioned between chloroform and water (50 mL each). The aqueous layer is again extracted with chloroform (25 mL) and the
15 combined organic layers dried (Na_2SO_4) and evaporated to leave a clear viscous oil which solidifies on standing. The solid is crystallized from ethyl acetate/hexanes as N-(4-carboxybenzyl)-N-methylnonanamide, m.p. 79.5-81°C.


20 EXAMPLE 16

N-((4-(1H-Tetrazol-5-yl)phenyl)methyl)nonanamide

To a solution of 4-(aminomethyl)benzonitrile (5.0 g, 0.038 moles) in 100 mL chloroform is added 3.82 g (0.038 moles) triethylamine. A solution of nonanoyl
25 chloride (6.68 g, 0.038 moles) in 10 mL chloroform is then added dropwise with stirring over a 10 min period and the final mixture stirred at room temperature for 18 hrs. The mixture is extracted with water (100 mL), saturated NaHCO_3 (50 mL), 2N HCl (50 mL), dried over
30 Na_2SO_4 and evaporated to leave 10.2 g crude N-(4-cyanobenzyl)nonanamide. This crude product is taken up in 50 mL dimethylformamide, to which is added 2.47 g (0.038 moles) sodium azide and 2.03 g (0.038 moles) ammonium chloride. The final mixture is heated at 90-
35 110°C for 4 hrs after cooling, the mixture is diluted with water (350 mL) and the resultant precipitate recovered by suction filtration, washed with water, and vacuum dried. Recrystallization from ethyl acetate left
40 N-[[4-(1H-tetrazol-5-yl)phenyl)methyl]nonanamide (2.6 g), m.p. 185-187°C.

CLAIMS (for contracting states BE, CH, DE, FR, GB,
IT, LI, LU, NL, SE)

1. A compound of the formula I:




wherein:


- 5 (a) R_1 is tetrazolyl or COOR' wherein R' is H or lower alkyl of 1 to 4 carbons, inclusive;
- (b) B is
- $$\begin{array}{c} \text{O} \\ || \\ -\text{C}- \end{array} \quad , \quad (\text{B}_1)$$

$$\begin{array}{c} \text{O} \\ || \\ -\text{S}- \end{array} \quad \text{or} \quad (\text{B}_2)$$


$$\begin{array}{c} \text{O} \\ || \\ \text{S} \\ || \\ -\text{C}-\text{NH}- \end{array} \quad (\text{B}_3);$$
- 10
- (c) X and Y are independently H or lower alkyl of 1 to 4 carbons, inclusive;
- (d) R_2 is alkylene, alkenylene, alkynylene branched or linear chains of 1 to 11 carbons, inclusive;
- 15 (e) Q is CH_3 , COOH , Br, NH_2 , H, imidazolyl, cyclohexyl,



5 or



- 10 wherein R₃ and R₄ are hydrogen or a C₁ to C₄ alkyl- or C₁ to C₄ alkoxy group with the proviso that when B is (B₁) and Q is H, R₂ is not methylene and with the further proviso that when B is (B₂) then Q cannot be CH₃, H or phenyl, or nontoxic, pharmaceutically acceptable acid addition or base salt thereof.
- 15 2. A compound according to claim 1 wherein R₁ is COOH having the formula:




wherein X, Y, B, R₂ and Q are as defined above.

3. A compound according to claim 2 wherein B is B₁.
4. A compound according to claim 3 wherein R₂ is a straight or branched alkylene chain of 3 to 11 carbons, inclusive.
- 20 5. A compound according to claim 3 wherein X is hydrogen or methyl and Y is hydrogen or methyl, and B, R₂ and Q are as defined above.

0226223

5 6. A compound according to claim 1 to 4, selected from
the group consisting of N-(4-carboxybenzyl)heptanamide,
N-(4-carboxybenzyl)nonanamide, N-(4-carboxybenzyl)
decanamide or the sodium salt thereof, N-(4-carboxy-
benzyl)octanamide, N-(4-carboxybenzyl)butyramide, N-
10 (4-carboxybenzyl)hexanamide, N-(4-carboxybenzyl)
undecanamide, N-1-[1-(4-carboxyphenyl)ethyl]nonanamide,
N-(4-carboxybenzyl)-N-methylnonamide.

7. A compound according to claim 3 wherein X, and Y
are each H, R_2 is alkylene or alkenylene branched or
15 straight chains of 1 to 3 carbons, inclusive, and Q is
(i)




20

wherein R_3 and R_4 are hydrogen or a C_1 to C_4 alkyl
or C_1 to C_4 alkoxy group

(ii) cyclohexyl or

25

(iii)



30

8. A compound according to claim 7, selected from the
group consisting of N-(4-carboxybenzyl)phenylacetamide,
N-(4-carboxybenzyl)-4-hydroxy-3-methoxycinnamamide, N-
35 (4-carboxybenzyl)cinnamamide, N-(4-carboxybenzyl)-4-
phenylbutyramide, N-(4-carboxybenzyl)cyclohexylacetamide
and N-(4-carboxybenzyl)-2-naphthoxyacetamide.

9. A compound of claim 1 wherein R_1 is tetrazolyl.


10. A compound of claim 9 and being N-[[4-(1H-tetrazol-5-yl)phenyl]methyl]nonanamide.

5 11. A process for the preparation of a compound of claim 1 to 10 which comprises contacting a compound of the formula




wherein R_1 , X and Y have the above mentioned meaning and a compound of the formula

15



20 wherein B, R_2 and Q have the above mentioned meaning and Hal is bromo, chloro or iodo to obtain a compound of formula I and, alternatively, further reacting the compound of formula I to obtain nontoxic pharmaceutically acceptable salts thereof.


25 12. A pharmaceutical composition comprising an effective amount of a compound of the formula I':



wherein:

35 (a) R_1 is tetrazolyl or COOR' wherein R' is H or lower alkyl of 1 to 4 carbons, inclusive;

(b) B is




0226223

5



or

10




(c) X and Y are independently H or lower alkyl of 1 to 4 carbons, inclusive;

15 (d) R_2 is alkylene, alkenylene, alkynylene branched or linear chains of 1 to 11 carbons, inclusive;

(f) Q is CH_3 , COOH , Br , NH_2 , H , imidazolyl, cyclohexyl,


20



25

or

30



wherein R_3 and R_4 are hydrogen or a C_1 to C_4 alkyl or a C_1 to C_4 alkoxy group

or nontoxic, pharmaceutically acceptable acid addition or base salt thereof; and a pharmaceutically acceptable carrier.


35

13. A method of using a compound of claim 12 for the manufacture of a medicament to reduce inflammation for treating pain or treating headaches in mammals.

0226223

CLAIMS (for contracting states AT, ES, GR)


1. A process for the manufacture of a compound of the formula I:



wherein:


- 5 (a) R_1 is tetrazolyl or COOR' wherein R' is H or lower alkyl of 1 to 4 carbons, inclusive;

- (b) B is



10

- (c) X and Y are independently H or lower alkyl of 1 to 4 carbons, inclusive;
- (d) R_2 is alkylene, alkenylene, alkynylene branched or linear chains of 1 to 11 carbons, inclusive;
- 15 (e) Q is CH_3 , COOH , Br, NH_2 , H, imidazolyl, cyclohexyl,




5 or



10 wherein R₃ and R₄ are hydrogen or a C₁ to C₄ alkyl- or C₁ to C₄ alkoxy group with the proviso that when B is (B₁) and Q is H, R₂ is not methylene and with the further proviso that when B is (B₂) then Q cannot be CH₃, H or phenyl, or nontoxic, pharmaceutically acceptable acid addition or base salt thereof,

15 which comprises contacting a compound of the formula



20


wherein R₁, X and Y have the above mentioned meaning and a compound of the formula

25 Hal-B-R₂-Q

wherein B, R₂ and Q have the above mentioned meaning and Hal is bromo, chloro or iodo to obtain a compound of formula I and, alternatively, further reacting the compound of formula I to obtain nontoxic pharmaceutically acceptable salts thereof.

30

2. A process according to claim 1 for manufacturing a compound wherein R₁ is COOH representing the formula:



35

wherein X, Y, B, R_2 and Q are as defined above.


3. A process according to claim 2 for preparing a compound wherein B is B_1 .

5 4. A process according to claim 3 for preparing a compound wherein R_2 is a straight or branched alkylene chain of 3 to 11 carbons, inclusive.

0 5. A process according to claim 3 for preparing a compound wherein X is hydrogen or methyl and Y is hydrogen or methyl, and B, R_2 and Q are as defined above.


15 6. A process according to claim 1 to 4 for preparing a compound selected from the group consisting of N-(4-carboxybenzyl)heptanamide, N-(4-carboxybenzyl)nonanamide, N-(4-carboxybenzyl)decanamide or the sodium salt thereof, N-(4-carboxybenzyl)octanamide, N-(4-carboxybenzyl)butyramide, N-(4-carboxybenzyl)hexanamide, N-(4-carboxybenzyl)undecanamide, N-1-[1-(4-carboxyphenyl)ethyl]nonanamide, N-(4-carboxybenzyl)-N-methylnonamide.

20 7. A process according to claim 3 for preparing a compound wherein X, and Y are each H, R_2 is alkylene or alkenylene branched or straight chains of 1 to 3 carbons, inclusive, and Q is
(i)



30 wherein R_3 and R_4 are hydrogen or a C_1 to C_4 alkyl or C_1 to C_4 alkoxy group
(ii) cyclohexyl or

35 (iii)




0226223

8. A process according to claim 7 for preparing a compound selected from the group consisting of N-(4-carboxybenzyl)phenylacetamide, N-(4-carboxybenzyl)-4-hydroxy-3-methoxycinnamamide, N-(4-carboxybenzyl)cinnamamide, N-(4-carboxybenzyl)-4-phenylbutyramide, N-(4-carboxybenzyl)cyclohexylacetamide and N-(4-carboxybenzyl)-2-naphthoxyacetamide.

9. A process of claim 1 for preparing a compound wherein R_1 is tetrazolyl.

10. A process of claim 9 for preparing N-[[4-(1H-tetrazol-5-yl)phenyl] methyl]nonanamide.


11. A method of using a compound of the formula I':




wherein:

(a) R_1 is tetrazolyl or COOR' wherein R' is H or lower alkyl of 1 to 4 carbons, inclusive;

(b) B is




or



(c) X and Y are independently H or lower alkyl of 1 to 4 carbons, inclusive;

- (d) R_2 is alkylene, alkenylene, alkynylene branched or linear chains of 1 to 11 carbons, inclusive;
(f) Q is CH_3 , $COOH$, Br, NH_2 , H, imidazolyl, cyclohexyl,

5



10

or

15



20

wherein R_3 and R_4 are hydrogen or a C_1 to C_4 alkyl or a C_1 to C_4 alkoxy group or nontoxic, pharmaceutically acceptable acid addition or base salt thereof; and a pharmaceutically acceptable carrier for the manufacture of a medicament to reduce inflammation for treating pain or treating headaches in mammals.

19



Europäisches Patentamt
European Patent Office
Office européen des brevets

11 Publication number:

0 226 223
A3

12

EUROPEAN PATENT APPLICATION

21 Application number: 86117748.3

22 Date of filing: 19.12.86

51 Int. Cl.: **C 07 C 103/46, C 07 C 103/84,
C 07 C 103/737, C 07 C 147/00,
C 07 C 157/07, C 07 D 257/04,
A 61 K 31/165, A 61 K 31/41**

30 Priority: 20.12.85 US 811567

43 Date of publication of application: 24.06.87
Bulletin 87/26

84 Designated Contracting States: AT BE CH DE ES FR GB
GR IT LI LU NL SE

88 Date of deferred publication of search
report: 03.01.90 Bulletin 90/1

71 Applicant: **WARNER-LAMBERT COMPANY, 201 Tabor
Road, Morris Plains New Jersey 07950 (US)**

72 Inventor: **Rafferty, Michael F., 59 Clear Lake Manor Drive
East, North Branford Connecticut 06471 (US)**
Inventor: **Johnson, Graham, 1130 Bandera Drive, Ann
Arbor Michigan 48103 (US)**

74 Representative: **Mansmann, Ivo, c/o Gödecke AG -
Patentabteilung Postfach 569 - Mooswaldallee 1-9,
D-7800 Freiburg (DE)**

54 **Benzoic acid and benzoic acid ester derivatives having anti-inflammatory and analgesic activity.**

57 Novel benzoic acid or benzoic acid ester derivatives, pharmaceutical compositions and methods of use thereof are the present invention. Utility is for the treatment of arthritis, asthma, Raynaud's disease, inflammatory bowel disorders, trigeminal or herpetic neuralgia, inflammatory eye disorders, psoriasis, dental pain, and headaches, particularly vascular headache, such as migraine, cluster, mixed vascular syndromes, as well as nonvascular, tension headaches.

EP 0 226 223 A3



European Patent
Office

EUROPEAN SEARCH REPORT

0226223

Application Number

EP 86 11 7748

DOCUMENTS CONSIDERED TO BE RELEVANT			
Category	Citation of document with indication, where appropriate, of relevant passages	Relevant to claim	CLASSIFICATION OF THE APPLICATION (Int. Cl. 4)
X	CHEMICAL ABSTRACTS vol. 91, no. 5, 30 July 1979, abstract no. 32654m, Columbus, Ohio, USA; J. A. MONTGOMERY et al.: "Analog of methotrexate" & J. Med. Chem. 1979, vol. 22, no. 7, pages 862-868 ---	1	C 07 C 103/46 C 07 D 257/04 C 07 C 103/84 C 07 C 103/737 C 07 C 143/75 C 07 C 157/07 A 61 K 31/165 A 61 K 31/41
X	CHEMICAL ABSTRACTS vol. 90, no. 25, 18 June 1979, abstract no. 203859t, Columbus, Ohio, USA; & JP - A- 5409266 (Y. TAKATORI) 24.01.1979 ---	1	
X,D	EP-A-0 031 954 (BOEHRINGER MANNHEIM GMBH) * claim 1 *; & US - A - 4443477 ---	1	
A,D	US-A-4 183 954 (A. BERTELLI) * claim 1 * ---	1	
A	AU-B- 492 025 (RORER ITALIANA S.P.A.) * claim 1 * ---	1	
A,D	EP-A-0 149 544 (PROCTER & GAMBLE COMPANY) * claim 1 * -----	1	
The present search report has been drawn up for all claims			TECHNICAL FIELDS SEARCHED (Int. Cl.4)
			C 07 C 103/00 C 07 D 257/00 C 07 C 157/00 C 07 C 143/00
Place of search		Date of completion of the search	Examiner
BERLIN		21-09-1989	PROBERT C.L.
CATEGORY OF CITED DOCUMENTS			
X : particularly relevant if taken alone Y : particularly relevant if combined with another document of the same category A : technological background O : non-written disclosure P : intermediate document T : theory or principle underlying the invention E : earlier patent document, but published on, or after the filing date D : document cited in the application L : document cited for other reasons ----- & : member of the same patent family, corresponding document			

**This Page is Inserted by IFW Indexing and Scanning
Operations and is not part of the Official Record**

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

- ☐ BLACK BORDERS
- ☐ IMAGE CUT OFF AT TOP, BOTTOM OR SIDES
- ☐ FADED TEXT OR DRAWING
- ☐ BLURRED OR ILLEGIBLE TEXT OR DRAWING
- ☐ SKEWED/SLANTED IMAGES
- ☐ COLOR OR BLACK AND WHITE PHOTOGRAPHS
- ☒ GRAY SCALE DOCUMENTS
- ☒ LINES OR MARKS ON ORIGINAL DOCUMENT
- ☐ REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY
- ☐ OTHER: _____

IMAGES ARE BEST AVAILABLE COPY.

As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.